

Sven Horvath, NGDC	Statistical	4.59					This statistical model computes the probability that sea ice will be present in concentrations above 25% for each grid cell in NSIDC's polar stereographic projection. Yearly data from 1980 through the present are used in a Bayesian binomial linear regression. Predictions include mean-center (SE) surface air temperature and geopotential height at 500mb. May monthly mean surface air temperature and geopotential height at 500mb. June sea ice concentration, and a trend index. This model predicts a minimum September sea ice extent of 4.59 million square km occurring on September 12th. Sea ice concentration data was obtained from NSIDC's Sea Ice Index V3 (Data Set ID: Q02135), and the air temperature and geopotential height data was from NASA's MERRA2 dataset.	Yearly data from 1980 through the present are used in a binomial linear regression to predict the probability that sea ice concentration will be above 25%. Predictions are made every other day in 2018. To estimate total sea ice extent, grid cells with a percentage above a certain threshold (chosen from a drop-out cross-validation test) are multiplied by the total area of the dataset provided by NSIDC's polar stereographic dataset and then summed. This model predicts a minimum September sea ice extent of 4.59 million km ² occurring on September 4th. Sea ice concentration data was obtained from NSIDC's Sea Ice Index V3 (Data Set ID: Q02135), and the air temperature and geopotential height data was from NASA's MERRA2 dataset.	NSIDC's Sea Ice Index V3 (Data Set ID: Q02135) NASA's MERRA2 dataset	
International Arctic Research Center	Statistical	4.69	4.69			Upper: 5.076 million sq. km.; Lower: 4.309 million sq. km.	The range assessments represent 95th and 5th percentile confidence intervals.	Our statistical model uses the NCEP/NCAP (R1) Reanalysis data sets to develop analog methods of atmospheric variables that correlate with sea ice extent. The R1 data covers the time period of 1948-present. The model generates an estimated deviation from the 1979-2017 September sea ice extent based by identifying the top 10th percentile for a number of sea ice and atmospheric variables using the April through June time period of each year, and then follows the seasonal decline in ice through the September period. The variables used are: 1) sea level pressure, 2) 500 mb height, 3) 3-mb temperature, 4) 925 mb temperature, and 5) sea surface temperature. A composite forecast is developed from a regression-weighted model.	Our model assumes no a priori knowledge of the current extent of Arctic sea ice. However, rely on the NSIDC published monthly September sea ice extents to estimate the sea-ice trend and use the same linear trend that monthly extent graphics.	Our model does not utilize sea ice thickness.
Dimitri Kondrashov (CUA)	Statistical	4.67	0.51				This statistical model forecast is based on inverse modeling techniques applied to the regional Arctic Sea Ice Extent (SIE).	This statistical model forecast is based on Data-Adaptive Hermite Decomposition (DAHD) and Multiscale Smart-Landau Models (MSLM) inverse modeling techniques applied to the regional Arctic Sea Ice Extent (SIE) from Sea Ice Index Version 3 dataset. The daily SIE data were aggregated to provide weekly-sampled dataset over eight Arctic sectors. DAHD-MSLM predictive model has been derived from SIE anomalies with annual cycle removed. The MSLM model is initialized from latest SIE conditions (July 2018) by ensemble of stochastic realizations to provide probabilistic, regional Arctic forecasts in September, as well as pan-Arctic ones.	1. Kondrashov, D., M. D. Chelton, and M. Chini. 2018. Data-adaptive hermite decomposition and prediction of Arctic sea ice extent. <i>Dynamics and Statistics of the Climate System</i> , 3(1), 66-110.10.1093/dms/dy001. 2. Chelton, M. D., and D. Kondrashov. 2017. Data-adaptive hermite spectra and multiyear smart-landau models. <i>Chaos</i> , 27, 091102. doi:10.1063/1.4988400	Forecasts were initialized from the pre-operational US Navy Global Ocean Forecasting System (GOFS) 1.1 for the ocean and sea ice using the Navy Coupled Ocean Data Assimilation (NCODA) system. The sea ice model assimilated GOFS 1.1 and AMSR sea ice concentration products. Atmospheric initial conditions were from the operational Navy Global Environmental Model (NAVMEM) using the Naval Atmospheric Variational Data Assimilation System (NAVDAS-AS).
NEL-NSM	Dynamic Model	4.7	31.3	0.81		4.3 to 5.2 Mkm ²	The uncertainty estimate is the range of the 10 member ensemble, and does not represent a full measure of uncertainty.	We performed ensemble forecasts with the Navy Earth System Modeling initial conditions on 2018-06-01 12Z through 2018-06-11 12Z. The atmospheric conditions were from NAVMEM-DOE v2.0. The sea ice model was part of the NAVMEM program at a 20km operational scale. The ocean and ice components were from the NSIDC sea ice model (NSIDC-SIEM) (Cummings 2005), which is a component of GOS 3.1 using HCEM and CICE (Cummings et al., 2016). SIE and SIE concentration are assimilated with NCODA (Poley et al., 2015). There was no bias correction performed on the results.	The ensemble forecasts were initialized using the observations from the GOFS 1.1 reinter files on the appropriate start date. Sea thickness products are not assimilated by GOFS 1.1.	
COM (D. Schroeder, et al.)	Statistical	4.7				< 0.5 mill. km ²	The given uncertainty is the mean forecast error based on forecasts for the years 1984 to 2017.	Based on May and June melt pond fraction we predict a mean 2018 SIE of 4.7 million km ² . This is slightly above the linear trend line, but lower than our previous SIE of 4.7 to 5.3 million km ² . While melt pond fraction has been generally low in May, the area covered by melt ponds in June is lower than the last 10 years in the Central Arctic and below in the Siberian part.	This is a statistical prediction based on the correlation between the ice area covered by melt ponds in May and ice extent in September. The melt pond area is derived from a simulation with the sea ice model CICE in which we incorporated a physics-based melt-pond model. See our publication in <i>Nature Climate Change</i> . http://www.nature.com/articles/nclimate2018018	include source (e.g., which data center), name (algorithm), DOI and/or data set website, and date (e.g., "NSIDC NASA Team, http://nsidc.org/data/seaice/DOE").
Ulfarok Kimura et al.	Statistical	4.71					Monthly mean ice extent in September will be about 4.7 million square kilometers. Our estimate is based on a statistical way using data from satellite microwave sensors. We used the sea ice thickness in December and ice movement from December to April. Predicted ice concentration map from July to September is available on our website: http://icr.arizona.edu/~ulfarok/ . Ice ice cover in the Laptev and East Siberian Seas will retreat with nearly same speed as last year. Minimum sea ice cover in September of this year will be the same as last year.	We predicted the Arctic sea-ice cover from coming July to November 1, using the data from satellite microwave sensor, AMSR-E (200203-201212) and AMSR2 (201212-201718). The analysis method is based on our recent research (Kimura et al., 2013). First, we expect the ice thickness distribution in April 30 from redistribution (divergence/convergence) of sea ice during December and April, based on the daily ice velocity data. Then, we predict the ice area based on the assumption that the ice thickness remains later and this ice melts sooner than the average. For the melt ponds, we used the partition homogeneity over the Arctic sea ice on December 15, traced the trajectories of the particles to the end of April using the satellite derived ice velocity (Kimura Dataset). Based on the relationship between particle density on April 15 and April concentration in summer, we predicted the summer sea ice cover of this year. We also take it into account that thickness of sea ice on December 1 calculated by an algorithm of Kirchfeld et al. (2014).	SIT dataset distributed by distributed by Arctic Data archive System (ADS). SIT dataset distributed by distributed by Arctic Data archive System (ADS). https://ads.nga.gov/ads/index.html ADS (AMER) years. SIT is calculated by an algorithm of Kirchfeld et al. (2014).	
Jianli Zhang and Axel Schweiger	Dynamic Model	4.72					Driven by the NCEP CFRS forecast atmospheric forcing, POMAAS is used to predict the September 2017 Arctic sea ice extent as well as ice thickness field and edge location, starting on July 1. The predicted September ice extent is 4.72 million square kilometers. The predicted ice thickness, fields and ice edge locations for September 2016 are also presented.	These results are obtained from a numerical seasonal forecasting system. The forecasting system is based on a synthesis of POMAAS, the NCEP CFRS forecast atmospheric forcing, and satellite observations of ice concentration. The CFRS forecast ranges from four to six months; there are a total of 58 CFRS ensemble forecasts run every two months, which four ensemble runs go out to 3 months, three runs go out to 2 months, and one run goes out to 45-day Gabe et al. (2016). These ensemble runs are used to create 6-hourly forecast atmospheric data that are widely accessible in real time, thus ideal for forecasting POMAAS forecasts. Data to seasonal time scales. We use a total of four CFRS forecast ensemble members from the POMAAS sea-ice ensemble forecasts. Ensemble member values from these four members are considered to be the prediction. To obtain the "best possible" initial conditions for the forecasts, we conducted a retrospective simulation that assimilates satellite ice concentration and sea surface temperature data through the end of May 2018 using the CFRS hindcast forcing data. After that, four ensemble POMAAS forecasts were conducted using atmospheric forecast forcing from four CFRS ensemble runs. Additional information about POMAAS prediction can be found in Zhang et al. (2018), Gabe, S., and others. The NCEP climate forecast system version 2.1. <i>Climatic</i> , 27, 2185-2208, 2014. Zhang, J., and D.A. Rothrock. Modeling global sea ice with thickness and ambly distribution model in general circulation coordinates. <i>Mon. Wea. Rev.</i> , 131(5), 681-697, 2003. Zhang, J., M. Swales, R. M. Lindsay, A. Schweiger, and J. Morrison. Ensemble one-year predictions of arctic sea ice for the spring and summer of 2008.	include source (e.g., which data center), name (algorithm), DOI and/or data set website, and date (e.g., "NSIDC NASA Team, http://nsidc.org/data/seaice/DOE").	
RASM (Kamal et al.)	Dynamic Model	4.75	0.468			0.239 million square kilometers	The uncertainty was estimated based on standard deviation of the 20-member ensemble.	We used RASM version 2.02, which is a recent version of the limited-area, fully coupled climate model consisting of the Weather Research and Forecasting (WRF), the Kansas National Laboratory/Atmospheric Ocean Program (KOP) and Sea Ice Model (SICM), Variable Resolution Coupled (VRC) hydrology and routing scheme (Majumdar et al., 2012; Roberts et al., 2012; D'Avolio et al., 2015; Hamman et al. 2016; Hamman et al., 2017; Cantua et al., 2017). The model uses CFRS as the initial and lateral boundary conditions and for nudging winds and temperature starting about 500 mb. We used one root case utilizing WRF37, including the Grid-3D parameterization scheme (with shallow cumulus convection option only turned on over the ocean grid).	For the July forecast we used one root case forced with CFRS to generate the initial conditions for the 20-member ensemble starting at 0000 UTC on July 1, 2018. The root case is a hindcast forecast from September 1979 through the end of June 2018, providing interannual and physically consistent initial conditions for the ensemble forecast using respective 20-member ensembles were initialized for a month forward using observational data from CFRS2. The CFRS2 forcing streams used for the ensemble were initialized everywhere (at 0000) between June 30 and July 1, 2018, and used for RASM forcing at time 0000 on July 1st 2018.	Self-generated from a hindcast run.
NCEP DOE	Dynamic Model	4.77	0.9			0.19	The standard deviation is calculated from the 20-member ensemble.	The outlook is produced from the Climate Prediction Center Experimental sea ice forecast system (EFSIS). The forecast is initialized from the Climate Forecast System Reanalysis (CFSR) for the ocean, land, and atmosphere and from the CPC sea ice initialization system (CSIS) for sea ice. Twenty forecast members are produced. Model bias that is removed is calculated based on 2006-2012 retrospective forecasts and corresponding observations.	Both sea ice concentration and sea ice thickness are initialized from the CPC sea ice initialization system (CSIS). The CSIS analysis is produced with the CSIS MOSM which uses surface fields from CFSR and assimilates satellite sea ice concentration retrieval from NSIDC NASA Team.	Both sea ice concentration and sea ice thickness are initialized from the CPC sea ice initialization system (CSIS). The CSIS analysis is produced with the CSIS MOSM which uses surface fields from CFSR and assimilates satellite sea ice concentration retrieval from NSIDC NASA Team.
Xiaojun Yuan, LEO	Statistical	4.78	18.95	0.6			The uncertainty of SIC prediction measured by RMSE was estimated based on cross-validated model experiments for 14 years.	The Linear Markov model has been developed to predict sea ice concentration to the pan-Arctic region of the seasonal time scale. The model employs 6 variables: NASA Team sea ice concentration, sea surface temperature (SST), surface air temperature, GH000, sector winds at GH000 (NCEP/NCAR reanalysis) for the period of 1979 to 2012. It is built in Markov process to predict the principal components forward one month at a time. The pan-Arctic sea ice concentration is calculated by summing up all of areas where predicted sea ice concentration exceeds 5%. Bias corrections have been applied to ice concentration predictions at grid points as well as the total sea ice extent prediction. The predictive skill of the model was evaluated by anomaly correlation between predictions and observations, and root-mean-square error (RMSE) in a false one-year cross-validated fashion. On average, the model is superior to the predictions by anomaly correlations, dated anomaly anomalies, and climatology (Yuan et al., 2016). For the three-month lead prediction of September sea ice concentration, the model has the higher skill (anomaly correlation) and lower RMSE in the Chukchi Sea and the Beaufort Sea than in other regions (Figure 4). The skill of the three-month lead prediction of the pan-Arctic sea ice extent in September is 0.88 with an RMSE of 0.46 million square kilometers. The Alaska regional SIC prediction is produced by a regional Linear Markov model developed by using SIC, SST, and SST in the EOF space following the NSIDC regional mask. The Alaska SIC forecast is calculated from predicted SIC. The skill of the regional SIC is 0.90 (correlation using cross-validated experiment) with RMSE of 0.22 million square kilometers. A similar model is used for Antarctic SIC forecast (Yuan and Yuan 2004).	NSIDC NASA Team. http://nsidc.org/data/seaice/DOE http://nsidc.org/data/seaice/DOE/ERA-interim-atmospheric-variables http://nsidc.org/data/seaice/DOE/NOAA-NCEP-ERSST-vert-sfc surface temperature data, http://nsidc.org/data/seaice/DOE/SFURF http://nsidc.org/data/seaice/DOE/SFURF http://nsidc.org/data/seaice/DOE/SFURF http://nsidc.org/data/seaice/DOE/SFURF	
Fengyan Wu and Robert Grumbine	Dynamic Model	4.84	10.62				The predicted Arctic minimum sea ice extent from the NCEP CFRS2 model with initial CFRS2 May and June initial conditions using the NCEP CFRS2 ensemble forecast is 4.84 million square kilometers with a standard deviation of 0.79 million square kilometers. The corresponding number for the Arctic's 1982 minimum square kilometers with a standard deviation of 0.63 million square kilometers.	We ran the NCEP CFRS2 model with 65 case of May and June 2018 and used initial conditions (SIC) that was modified from the initial CFRS2 data of each day at 00Z by thinning the Arctic ice pack based on bias from previous year sea ice outside. If this thinning would have eliminated ice from areas observed to have sea ice, a minimum thickness of 0.1 m was left in place for the ice. Bias correction was applied to the Arctic sea ice extent.	Include source (e.g., which data center), name (algorithm), DOI and/or data set website, and date (e.g., "NSIDC NASA Team, http://nsidc.org/data/seaice/DOE"). http://nsidc.org/data/seaice/DOE http://nsidc.org/data/seaice/DOE http://nsidc.org/data/seaice/DOE http://nsidc.org/data/seaice/DOE	NCEP CFRS2 model goals with bias correction for the Arctic (May 1-June 30, 2018)

FD-ESM (Qian et al.)	Dynamic Model	Coupled dynamical models	5.2							Our prediction is based on FD-ESM (the First Institute of Oceanography-Berth System Model) with data assimilation. The prediction of September pan-Arctic extent in 2018 is 5.2 (±0.3) million square kilometers. 5.2 and 6.5 million square kilometers is the average and one standard deviation of 10 ensemble members, respectively.	This is a model contribution. The initialization is also from the same model (FD-ESM) but with data assimilation. The data assimilation method is Ensemble Adjustment Kalman Filter (EAKF). The data of SST (sea surface temperature) and SLA (sea level anomaly) from 1 January 1992 to 1 July 2018 are assimilated into FD-ESM model to get the initial condition for the prediction of the Arctic Sea ice. There is no sea ice data assimilation.	No dataset are used for initial sea ice concentration.	No dataset are used for initial sea ice thickness.
AWI consortium (Kauker et al.)	Dynamic Model	Ocean-sea ice dynamical models	5.2	0.14	Ensemble spread of the forcing years 2008 to 2017 used by the sea ice - ocean model from Apr 8 to end of September).					For the present outlook the coupled ice-ocean model (MOCIM) has been forced with atmospheric surface data from January 1948 to July 8th 2018 (combination of NCEP/NCAR and NCEP-CFSR and NCEP-CFSv2). All ensemble model experiments have been started from the same initial conditions on July 8th 2018. The model setup has not changed with respect to the IGO in 2015. We used atmospheric forcing data from each of the years 2008 to 2017 for the ensemble prediction and thus obtain 10 different realizations of potential sea ice evolution for the summer of 2018. The use of an ensemble allows to estimate probabilities of sea-ice extent predictions for September 2018. A variational assimilation system (MOCIM) has been used to initialize the model using the Alfred Wegener Institute's CryoSat-2 ice thickness product, the University of Bremen's snow depth product, and the OSI SAF ice concentration and sea-surface temperature products. Observations from March and April were used. A bias correction scheme for the CryoSat-2 ice thickness which employs a spatially variable scaling factor could enhance the skill considerably (Kauker et al., 2015; http://www.the-cryosphere-discuss.net/9-2015-117/).	OSI SAF EUMETSAT OSI-401 March and April 2018	CryoSat-2 from Alfred Wegener Institute of March and April 2018	